

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec[®]

EPA Reg. No. 62719-375

Control of Powdery Mildew in Melons[†]

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec[®] fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Directions for Use

[†]Melons including, but not limited to: Citron Melon, Muskmelon (Cantaloupe, Casaba, Crenshaw Melon, Golden Pershaw Melon, Honeydew Melon, Honey Balls, Mango Melon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon, True Cantaloupe), Watermelon

Disease	Recommended Spray Interval (days)	Rate of Quintec (fl oz/acre)
powdery mildew (<i>Sphaerotheca fuliginea</i>)	10 - 14	4 - 6

Apply uniformly in a spray volume that provides thorough coverage of the crop. A minimum spray volume of 30 gallons per acre is recommended. Disease control may be reduced at low spray volumes or if spray coverage is inadequate.

Application Timing: Apply Quintec before visible symptoms of powdery mildew appear. Quintec will not control latent or established infections of powdery mildew. If powdery mildew infection is established, Quintec should be applied in tank mix combination with a curative fungicide or following application of a curative fungicide. If other diseases are present, use fungicides that are effective on those diseases. If reliable predictive modeling (risk index) systems are available and indicate low disease pressure conditions, minimum label rates and longer application intervals may be used. Maximum label rates and short application intervals are recommended for severe or threatening disease conditions.

Use Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- **Maximum Application Rate:** Do not apply more than 6 fl oz of Quintec per acre per application or more than 24 fl oz of Quintec per acre per crop.
- Do not make more than 4 applications of Quintec per crop.
- It is recommended to alternate use of Quintec with a product with a different mode of action after each application of Quintec.
- Do not make more than 2 consecutive applications of Quintec before rotating to a product that is effective on powdery mildew and has a different mode of action than Quintec. In either case, apply the alternate product within 14 days of applying Quintec.

- Carefully inspect the crop several days after each application of Quintec and at regular intervals during crop development. Under certain environmental conditions, Quintec may cause leaf spotting and chlorosis. If these symptoms occur after applying Quintec, discontinue its use.
- Do not aerially apply this product.
- Do not graze or harvest forage from cover crops treated with Quintec.
- **Chemigation:** Do not apply this product through any type of irrigation system.
- Carefully read and follow all applicable use directions, precautions, and limitations on the respective product labels used in mixtures.

Rotational Crop Restrictions: Crops grown for food or feed without registered uses for quinoxifen may be replanted 12 months after the last application of quinoxifen.

Note: This product is toxic to fish and aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

®Trademark of Dow AgroSciences LLC

R136-008

Approved: 09/11/06

Initial printing.